

THE CHEMMACROS BUNDLE

v4.2a 2013/11/04

packages **CHEMMACROS** (v4.2b), **CHEMFORMULA** (v4.2a), **GHSYSTEM** (v4.0) and
CHEMGREEK (v0.2)

documentation for the **CHEMFORMULA** package

Clemens NIEDERBERGER

<https://bitbucket.org/cgnieder/chemmacros/>

contact@mychemistry.eu




Table of Contents

| | | | | | |
|----------|---------------------------------|----------|--------------|---------------------------------------|-----------|
| 1 | Introduction | 2 | 7.4 | Charges and Other Superscripts | 8 |
| | | | 7.5 | Bonds | 10 |
| 2 | News | 2 | 7.5.1 | Native Bonds | 10 |
| 2.1 | Version 4.0 | 2 | 7.5.2 | Flexible Bonds | 10 |
| 2.2 | Version 4.2 | 2 | 7.6 | Customization | 12 |
| 3 | Licence and Requirements | 3 | 7.7 | Standalone Formulae | 15 |
| 4 | Setup | 3 | 8 | Special Input Types | 15 |
| 5 | The Basic Principle | 3 | 8.1 | Single Token Groups | 16 |
| 6 | Stoichiometric Factors | 5 | 8.2 | Option Input | 16 |
| 7 | Compounds | 7 | 9 | Escaped Input | 17 |
| 7.1 | Adducts | 7 | 9.1 | Text | 17 |
| 7.2 | Subscripts | 8 | 9.2 | Math | 18 |
| 7.3 | Commands | 8 | 10 | Arrows | 18 |
| | | | 10.1 | Arrow types | 18 |
| | | | 10.2 | Labels | 20 |

| | | | |
|-----------------------------------|----|--|----|
| 10.3 Customization | 21 | 13 Usage In Math Equations | 27 |
| 10.4 Modify Arrow Types | 22 | 14 Usage with TikZ or pgfplots and externalization | 28 |
| 10.5 Standalone Arrows | 24 | 15 Lewis Formulae | 29 |
| 11 Names | 24 | 16 Further Examples | 30 |
| 11.1 Syntax | 24 | References | 33 |
| 11.2 Customization | 25 | Index | 34 |
| 11.3 Standalone Names | 25 | | |
| 12 Format and Font | 25 | | |

1 Introduction

Probably every chemist using \LaTeX $_{2\epsilon}$ is aware of the great mhchem package by Martin Hensel. There have always been some difficulties intertwining it with **CHEMMACROS**, though. Also, some other minor points in mhchem always bothered me, but they hardly seemed enough for a new package. They weren't even enough for a feature request to the mhchem author. The challenge and the fun of creating a new package and the wish for a highly customizable alternative led to **CHEMFORMULA** after all.

CHEMFORMULA works very similar to mhchem but is more strict as to how compounds, stoichiometric factors and arrows are input. In the same time **CHEMFORMULA** offers *many* possibilities to customize the output.

2 News

2.1 Version 4.0

Introduced in
version 4.0

Since version 4.0, the **CHEMFORMULA** package can be used independently from **CHEMMACROS**. This means that if you say

```
\usepackage{chemformula}
```

then **CHEMMACROS** will not be loaded. The **CHEMMACROS** package, however, *will* load **CHEMFORMULA**.

2.2 Version 4.2

- New option **arrow-style**.
- New command **\chlewis** that allows to add Lewis electrons to an atom, see section 15.

3 Licence and Requirements

Permission is granted to copy, distribute and/or modify this software under the terms of the L^AT_EX Project Public License (LPPL), version 1.3 or later (<http://www.latex-project.org/lppl.txt>). The software has the status “maintained.”

The **CHEMFORMULA** package needs and thus loads the packages l3kernel [The13a], xparse, l3keys2e and xfrac (all three are part of the l3packages bundle [The13b]), tikz¹ [Tan10], ams-math [Ameo2], nicefrac [Rei98] and scrfile (from the KOMA-Script² bundle [KN12]).

4 Setup

All of **CHEMFORMULA**’s options belong to **CHEMMACROS**’ module `chemformula`. This means they can be setup with

```
\chemsetup[chemformula]{<options>}
```

Set up options for **CHEMFORMULA** exclusively, or

```
\chemsetup{chemformula/<option1>,chemformula/<option2>}
```

Set up options for **CHEMFORMULA** together with others of **CHEMMACROS**’ options.

However, if you’re using **CHEMFORMULA** as a standalone package the command `\chemsetup` is not available. This is why **CHEMFORMULA** also has its own setup command:

```
\setchemformula{<options>}
```

Set up **CHEMFORMULA** when using it independently from **CHEMMACROS**.

5 The Basic Principle

CHEMFORMULA offers one main command.

```
\ch[<options>]{<input>}
```

CHEMFORMULA’s main command.

The usage will seem very familiar to you if you’re familiar with mhchem:

1. on CTAN as pgf: <http://mirrors.ctan.org/graphics/pgf/>

2. on CTAN as koma-script: <http://mirrors.ctan.org/macros/latex/contrib/koma-script/>

| | |
|---|---|
| 1 <code>\ch{H2O}</code> <code>\par</code> | H_2O |
| 2 <code>\ch{Sb2O3}</code> <code>\par</code> | Sb_2O_3 |
| 3 <code>\ch{H+}</code> <code>\par</code> | H^+ |
| 4 <code>\ch{CrO4^2-}</code> <code>\par</code> | CrO_4^{2-} |
| 5 <code>\ch{AgCl2-}</code> <code>\par</code> | AgCl_2^- |
| 6 <code>\ch{[AgCl2]-}</code> <code>\par</code> | $[\text{AgCl}_2]^-$ |
| 7 <code>\ch{Y^{99+}}</code> <code>\par</code> | Y^{99+} |
| 8 <code>\ch{Y^{99+}}</code> <code>\par</code> | Y^{99+} |
| 9 <code>\ch{H2_{(aq)}}</code> <code>\par</code> | $\text{H}_{2(\text{aq})}$ |
| 10 <code>\ch{NO3-}</code> <code>\par</code> | NO_3^- |
| 11 <code>\ch{(NH4)2S}</code> <code>\par</code> | $(\text{NH}_4)_2\text{S}$ |
| 12 <code>\ch{^{227}_{90}Th+}</code> <code>\par</code> | $^{227}_{90}\text{Th}^+$ |
| 13 <code>\$V_{\ch{H2O}}\$</code> <code>\par</code> | $V_{\text{H}_2\text{O}}$ |
| 14 <code>\ch{Ce^{IV}}</code> <code>\par</code> | Ce^{IV} |
| 15 <code>\ch{KCr(SO4)2 * 12 H2O}</code> | $\text{KCr}(\text{SO}_4)_2 \cdot 12 \text{H}_2\text{O}$ |

However, there are differences. The most notable one: **CHEMFORMULA** distinguishes between different types of input. These different parts *have* to be separated with blanks:

`\ch{part1 part2 part3 part4}`

A blank in the input *never* is a blank in the output. This role of the blank strictly holds and disregarding it can have unexpected results and even lead to errors.

Another notable difference: **CHEMFORMULA** tries to avoid math mode whenever possible:

| | |
|--|---------------------------|
| 1 <code>\ch{A + B ->[a] C}</code> <code>\par</code> | $A + B \xrightarrow{a} C$ |
| 2 <code>\ce{A + B ->[a] C}</code> | $A + B \xrightarrow{a} C$ |

This means that `\ch{2H2O}` is recognized as a *single* part, which in this case is recognized as a compound.

| | |
|--|------------------------|
| 1 <code>\ch{2H2O}</code> <code>\par</code> | $2\text{H}_2\text{O}$ |
| 2 <code>\ch{2 H2O}</code> | $2 \text{H}_2\text{O}$ |

This also means, that a part cannot contain a blank since this will automatically divide it into two parts. If you need an extra blank in the output you need to use `~`. However, since commands in most cases gobble a space after them a input like `\ch{\command ABC}` will be treated as a single part. If you want or need to divide them you need to add an empty group: `\ch{\command{} ABC}`. The different input types are described in the following sections.

There are some options to customize the output of the `\ch` command. They can either be applied locally using the optional argument or can be set globally using the `setup` command. All options of **CHEMFORMULA** belong to the module `chemformula` and can be set in different ways:

`\chemsetup[chemformula]{<options>}`

when loaded via **CHEMMACROS**

`\chemsetup{chemformula/<options>}`

when loaded via **CHEMMACROS**

`\setchemformula{<options>}`

independent from **CHEMMACROS**

6 Stoichiometric Factors

A stoichiometric factor may only contain of numbers and the signs `.`, `-`, `/`, `()`

| | | | |
|----|-------------------------|-------------------|----------------|
| 1 | <code>\ch{2}</code> | <code>\par</code> | |
| 2 | <code>\ch{12}</code> | | |
| 3 | | | 2 |
| 4 | % decimals: | | 12 |
| 5 | <code>\ch{.5}</code> | <code>\par</code> | .5 |
| 6 | <code>\ch{5,75}</code> | | 5.75 |
| 7 | | | $\frac{3}{2}$ |
| 8 | % fractions: | | $1\frac{1}{2}$ |
| 9 | <code>\ch{3/2}</code> | <code>\par</code> | (1/2) |
| 10 | <code>\ch{1_1/2}</code> | | |
| 11 | | | |
| 12 | % "iupac": | | |
| 13 | <code>\ch{(1/2)}</code> | | |

As you can see if you input decimal numbers a missing leading zero is added.
You have to be a little bit careful with the right syntax but I believe it is rather intuitive.

1 this won't work but will result in an error: `\ch{1/1_1}`

If stoichiometric factors are enclosed with parentheses the fractions are not recognized and missing leading zeros are not added. What's inside the parentheses is typeset as is.

1 `\ch{(1/2) H2O}` `\ch{1/2 H2O}` `\ch{0.5 H2O}`

$(1/2)\text{H}_2\text{O}$ $\frac{1}{2}\text{H}_2\text{O}$ $0.5\text{H}_2\text{O}$

You can find many examples like the following for stoichiometric factors in parentheses in the IUPAC Green Book [Coh+08]:



There are a few possibilities to customize the output.

`decimal-marker = {\langle marker \rangle}` Default: `.`

The symbol to indicate the decimal.

`frac-style = math|xfrac|nicefrac` Default: `math`

Determines how fractions are displayed.

`frac-math-cmd = {\langle command sequence \rangle}` Default: `\frac`

Introduced in
version 4.1

Allows you to choose which command is used with `frac-style = math`. This needs to be a command sequence that takes two arguments that are set in math mode.

`stoich-space = {\langle skip \rangle}` Default: `.1667em plus .0333em minus .0117em`

The space that is placed after the stoichiometric factor. A rubber length.

`stoich-paren-parse = true|false` Default: `false`

If set to true stoichiometric factors enclosed by parentheses also are parsed.

```
1 \ch[decimal-marker={,}]{3.5} \ch[decimal-marker={\cdot}]{3,5}
```

3,5 3·5

The option `frac-style = xfrac` uses the `\sfrac` command of the `xfrac` package. The output strongly depends on the font you use.

```
1 \ch[frac-style=xfrac]{3/2} \ch[frac-style=xfrac]{1_1/2}
```

$\frac{3}{2}$ $1\frac{1}{2}$

CHEMFORMULA defines the instance `formula-text-fraction` which you can redefine to your needs. See the `xfrac` documentation for further information. The default definition is this:

```
1 \DeclareInstance{xfrac}{chemformula-text-fraction}{text}
2 {
3   slash-left-kern = -.15em ,
4   slash-right-kern = -.15em
5 }
```

This document uses the font Linux Libertine O and the following definition:

```
1 \DeclareInstance{xfrac}{chemformula-text-fraction}{text}
2 {
3   scale-factor      = 1 ,
4   denominator-bot-sep = -.2ex ,
```

7 Compounds

```
5 denominator-format = \scriptsize #1 ,
6 numerator-top-sep   = -.2ex ,
7 numerator-format     = \scriptsize #1 ,
8 slash-right-kern     = .05em ,
9 slash-left-kern      = .05em
10 }
```

The option `frac-style = nicefrac` uses the `\nicefrac` command of the `nicefrac` package.

```
1 \ch[frac-style=nicefrac]{3/2} \ch[frac-style=nicefrac]{1_1/2}
```

$\frac{3}{2}$ $1\frac{1}{2}$

The option `stoich-space` allows you to customize the space between stoichiometric factor and the group following after it.

```
1 \ch{2 H2O} \par 2 H2O
2 \ch[stoich-space=.3em]{2 H2O} 2 H2O
```

7 Compounds

CHEMFORMULA determines compounds as the type that “doesn’t fit in anywhere else.” This point will become more clear when you know what the other types are.

```
1 \ch{H2SO4} \par H2SO4
2 \ch{[Cu(NH3)4]^2+} [Cu(NH3)4]2+
```

7.1 Adducts

CHEMFORMULA has two identifiers which will create adducts.

```
\ch{A.B}
A·B
```

```
\ch{A*B}
A·B
```

```
1 \ch{CaSO4.H2O} \par CaSO4·H2O
2 \ch{CaSO4*H2O} CaSO4·H2O
```

Since numbers in a compound always are treated as subscripts (see section 7.2) you sometimes need to introduce stoichiometric factors for the right output:

| | |
|---|---|
| <ol style="list-style-type: none"> 1 <code>\ch{Na3PO4*12H2O} \par</code> 2 <code>\ch{Na3PO4* 12 H2O} \par</code> 3 <code>\ch{Na3PO4 * 12 H2O}</code> | $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ |
|---|---|

7.2 Subscripts

All numbers in a compound are treated as subscripts.

| | |
|---|-------------------------|
| <ol style="list-style-type: none"> 1 <code>\ch{H2SO4}</code> | H_2SO_4 |
|---|-------------------------|

If you want a letter to be a subscript you can use the math syntax:

| | |
|--|----------|
| <ol style="list-style-type: none"> 1 <code>\ch{A_nB_m}</code> | A_nB_m |
|--|----------|

The subscript recognizes groups. You can also use math inside it.

| | |
|--|--------------------------------------|
| <ol style="list-style-type: none"> 1 <code>\ch{A_{n\$}B_{m\$}} \par</code> 2 <code>\ch{NaCl_{(aq)}}</code> | A_nB_m $\text{NaCl}_{(\text{aq})}$ |
|--|--------------------------------------|

7.3 Commands

Commands are allowed in a compound:

| | |
|--|---|
| <ol style="list-style-type: none"> 1 <code>\ch{\textbf{A2}B3} \ch{A2\color{red}B3}</code> | $\text{A}_2\text{B}_3 \quad \text{A}_2\text{B}_3$ |
|--|---|

However, if the commands demand numbers as argument, *e. g.*, space commands or **CHEM-MACROS'** `\ox` command the direct use will fail. This is because the numbers are treated as subscripts *before* the command expands.

| |
|--|
| <ol style="list-style-type: none"> 1 <code>\ch{A\hspace{2mm}B}</code> will raise an error because <code>\hspace</code> sees something like 2 this: <code>\hspace{\$_2\$mm}</code>. Actually not at all like it but equally bad <code>\ldots</code> |
|--|

See section 9.1 for a way around this.

7.4 Charges and Other Superscripts

Basics If a compound *ends* with a plus or minus sign it will be treated as charge sign and typeset as superscript. In other places a plus is treated as a triple bond and a dash will be used as a single bond, see section 7.5.

7 Compounds

| | |
|---------------------------------------|-------------------|
| 1 <code>\ch{A+B} \ch{AB+} \par</code> | $A \equiv B AB^+$ |
| 2 <code>\ch{A-B} \ch{AB-}</code> | $A - B AB^-$ |

For longer charge groups or other superscripts you can use the math syntax. It recognizes groups and you can use math inside them. Inside these groups neither + nor - are treated as bonds. If a dot . is inside a superscript it is treated as indicator for a radical. A * gives the excited state.


| | |
|--|------------------|
| 1 <code>\ch{A^{x-}} \par</code> | A^{x-} |
| 2 <code>\ch{A^x-} \par</code> | A^{x-} |
| 3 <code>\ch{A^{x}-} \par</code> | A^{x-} |
| 4 <code>\ch{A^{\\$x-\\$}} \par</code> | A^{x-} |
| 5 <code>\ch{RNO2^{\cdot-}} \par</code> | $RNO_2^{\cdot-}$ |
| 6 <code>\ch{^3H} \par</code> | 3_1H |
| 7 <code>\ch{^{14}6C} \par</code> | $^{14}_6C$ |
| 8 <code>\ch{^{58}_{26}Fe} \par</code> | $^{58}_{26}Fe$ |
| 9 <code>\ch{NO^*}</code> | NO^* |

Ions and ion composites with more than one charge can be typeset quite as easy:

| | |
|--|-------------------------------|
| 1 <code>\ch{SO4^2-} \ch{Ca^2+ SO4^2-}</code> | $SO_4^{2-} Ca^{2+} SO_4^{2-}$ |
|--|-------------------------------|


Charge Commands You don't need to use `\mch` and related commands inside `\ch`. Indeed, you *shouldn't* use them as they might mess with the subscript and superscript alignment. The **CHEMMACROS** option circled is obeyed by `\ch`.

```
1 \chemsetup[option]{circled=all}
2 \ch{H+ + OH- <=> H2O}
```



Behaviour The superscripts behave differently depending on their position in a compound, if there are super- and subscripts following each other directly.

```
1 \ch{^33B} \ch{{}^33B} \ch{3^3B} \ch{B^3} \ch{B3^3} \par
2 \ch{^{23}_{123}B} \ch{{}^{23}_{123}B} \ch{{}_{123}^{23}B}
3 \ch{B^{23}} \ch{B_{123}^{23}} \par
4 \ch{^{123}_{23}B} \ch{{}^{123}_{23}B} \ch{{}_{23}^{123}B}
5 \ch{B^{123}} \ch{B23^{123}}
```



- If a compound *starts* with a sub- or superscript both sub- and superscript are aligned to the *right* else to the *left*.
- If a compound *does not start* with a sub- or superscript and there is both a sub- and a superscript, the superscript is shifted additionally by a length determined from the option `charge-hshift = {<dim>}`, also see page 12f.

The second point follows IUPAC's recommendations:

In writing the formula for a complex ion, spacing for charge number can be added (staggered arrangement), as well as parentheses: SO_4^{2-} , $(\text{SO}_4)^{2-}$. The staggered arrangement is now recommended.

IUPAC Green Book [Coh+08, p. 51]

7.5 Bonds

7.5.1 Native Bonds

There are three kinds of what I will call “native bonds”:

| | |
|---|------------------------------------|
| 1 single: <code>\ch{CH3-CH3}</code> <code>\par</code> | single: $\text{CH}_3\text{—CH}_3$ |
| 2 double: <code>\ch{CH2=CH2}</code> <code>\par</code> | double: $\text{CH}_2=\text{CH}_2$ |
| 3 triple: <code>\ch{CH+CH}</code> | triple: $\text{CH}\equiv\text{CH}$ |

7.5.2 Flexible Bonds

Predefined Bonds In addition to the three native bonds there are a few more which can be called by

`\bond{<bond name>}`

Prints the bond type specified by `<bond name>`.

The predefined bond types are shown in table 1 on the following page.

1 `\ch{C\bond{sb}C\bond{db}C\bond{tp}C\bond{deloc}C\bond{tde loc}C\bond{co>}C\bond{<co}C}`




TABLE 1: Bonds available with `\bond`.

| name | appearance | aliases |
|--------|------------|------------|
| single | — | normal, sb |
| double | = | db |
| triple | ≡ | tp |
| dotted | | semisingle |
| deloc | — — | semidouble |
| tdeloc | ≡ ≡ | semitriple |
| co> | → | coordright |
| <co | ← | coordleft |

Own Bonds `CHEMFORMULA` offers commands to define own bond types:

`\DeclareChemBond{<name>}{<code>}`

Define the new bond type `<name>`.

`\RenewChemBond{<name>}{<code>}`

Redefine the existing bond type `<name>`.

`\DeclareChemBondAlias{<new name>}{<old name>}`

Declare the bond type `<new name>` to be an alias of `<old name>`.

`\ShowChemBond{<name>}`

Print the definition of bond type `<name>`.

The usage is best described with an example. So let's see how the single bond and the `co>` bond are defined:

```

1 \DeclareChemBond{single}
2   { \draw[chembond] (chemformula-bond-start) -- (chemformula-bond-end) ; }
3 \DeclareChemBond{coordright}
4   {
5     \draw[chembond,butt cap->]
6       (chemformula-bond-start) -- (chemformula-bond-end) ;
7   }
8 \DeclareChemBondAlias{co>}{coordright}

```

Two points are important: the names of the starting and the ending coordinates, `chemformula-bond-start` and `chemformula-bond-end`, and the TikZ style of the bonds `chembond`.

So, let's say you want to define a special kind of dashed bond. You could do this:

```

1 \usetikzlibrary{decorations.pathreplacing}
2 \makeatletter
3 \DeclareChemBond{dashed}
4 {
5   \draw[
6     chembond,
7     decorate,
8     decoration={
9       ticks,
10      segment length=\chemformula@bondlength/10,amplitude=1.5pt
11    }]
12    (chemformula-bond-start) -- (chemformula-bond-end) ;
13 }
14 \makeatother
15 \chemsetup[chemformula]{bond-length=2ex}
16 \ch{C\bond{dashed}C}

```



The last example showed you another macro: `\chemformula@bondlength`. It only exists so you can use it to access the bond length as set with `bond-length` directly.

7.6 Customization

These options allow you to customize the output of the compounds:

| | |
|--|------------------|
| <code>subscript-vshift = {<dim>}</code> | Default: 0pt |
| Extra vertical shift of the subscripts. | |
| <code>subscript-style = text math</code> | Default: text |
| Style that is used to typeset the subscripts. | |
| <code>charge-hshift = {<dim>}</code> | Default: .25em |
| Shift of superscripts when following a subscript. | |
| <code>charge-vshift = {<dim>}</code> | Default: 0pt |
| Extra vertical shift of the superscripts. | |
| <code>charge-style = text math</code> | Default: text |
| Style that is used to typeset the superscripts. | |
| <code>adduct-space = {<dim>}</code> | Default: .1333em |
| Space to the left and the right of the adduct point. | |
| <code>bond-length = {<dim>}</code> | Default: .5833em |
| The length of the bonds. | |

bond-offset = $\{\langle dim \rangle\}$ Default: .07em
Space between bond and atoms.

bond-style = $\{\langle TikZ \rangle\}$ (initially empty)
TikZ options for the bonds.

bond-penalty = $\{\langle num \rangle\}$ Default: 10000
The penalty that is inserted after a bond for (dis-)allowing line breaks.

Introduced in
version 4.0a

radical-style = $\{\langle TikZ \rangle\}$ (initially empty)
TikZ options for the radical point.

radical-radius = $\{\langle dim \rangle\}$ Default: .2ex
The radius of the radical point.

radical-hshift = $\{\langle dim \rangle\}$ Default: .15em
Horizontal shift before the radical point is drawn.

radical-vshift = $\{\langle dim \rangle\}$ Default: .5ex
Vertical shift relative to the current baseline.

radical-space = $\{\langle dim \rangle\}$ Default: .15em
Horizontal shift after the radical point is drawn.

Maybe you have noticed that charges of certain ions are shifted to the right.

1 `\ch{SO4^2-} \ch{NH4+} \ch{Na+}` $\text{SO}_4^{2-} \text{NH}_4^+ \text{Na}^+$

They are shifted if they *follow* a subscript which follows IUPAC recommendations [Coh+08, p. 51]. The amount of the shift can be set with the option **charge-hshift** = $\{\langle . \rangle\}$

```
1 \ch{SO4^2-} \ch{NH4+} \ch{Na+} \par
2 \chemsetup[chemformula]{charge-hshift=.5ex}
3 \ch{SO4^2-} \ch{NH4+} \ch{Na+} \par
4 \chemsetup[chemformula]{charge-hshift=.5pt}
5 \ch{SO4^2-} \ch{NH4+} \ch{Na+}
```


$\text{SO}_4^{2-} \text{NH}_4^+ \text{Na}^+$
 $\text{SO}_4^{2-} \text{NH}_4^+ \text{Na}^+$
 $\text{SO}_4^{2-} \text{NH}_4^+ \text{Na}^+$

Despite IUPAC's recommendation **CHEMFORMULA** does not make fully staggered arrangements in the default setting as I find it hard to read in some cases and ugly in others. Since this is a subjective decision **CHEMFORMULA** not only let's you define the absolute amount of the shift but also provides a possibility for full staggered arrangements. For this you have to use **charge-hshift** = full.

```

1 \ch[charge-hshift=0pt]{C5H11+} \ch[charge-hshift=0pt]{SO4^2-} \par
2 \ch{C5H11+} \ch{SO4^2-} \par
3 \ch[charge-hshift=1ex]{C5H11+} \ch[charge-hshift=1ex]{SO4^2-} \par
4 \ch[charge-hshift=full]{C5H11+} \ch[charge-hshift=full]{SO4^2-}

```




If you don't want the charges to be typeset in text mode you can switch to math mode:

```

1 \ch{M^x+} \ch{SO4^2-} \par
2 \chemsetup[chemformula]{charge-style = math}
3 \ch{M^x+} \ch{SO4^2-}

```




The option `subscript-vshift` can be used to adjust the vertical shift of the subscripts:

```

1 \ch{H2SO4} \ch{Na3PO4} \par
2 \chemsetup[chemformula]{subscript-vshift=.5ex}
3 \ch{H2SO4} \ch{Na3PO4} \par
4 \chemsetup[chemformula]{subscript-vshift=-.2ex}
5 \ch{H2SO4} \ch{Na3PO4}

```





You can choose the mode subscripts are typeset in the same way as it is possible for the charges:

```

1 \ch{A_nB_m} \ch{H2SO4} \par
2 \chemsetup[chemformula]{subscript-style = math}
3 \ch{A_nB_m} \ch{H2SO4}


```






The option `adduct-space` sets the space left and right to the adduct symbol `.`.

```
1 \ch{Na3PO3*H2O} \par
2 \chemsetup[chemformula]{adduct-space=.2em}
3 \ch{Na3PO3*H2O}
```




Changing the length of the bonds:

```
1 \chemsetup[chemformula]{bond-length=4mm}%
2 single: \ch{CH3-CH3} \par
3 double: \ch{CH2=CH2} \par
4 triple: \ch{CH+CH}
```



You can change the distance between bond and atom, too:

```
1 \ch{H-H + N+N + O=O} \par
2 \ch[bond-offset=1pt]{H-H + N+N + O=O}
```



7.7 Standalone Formulae

Introduced in
version 4.0

CHEMFORMULA offers a command that *only accepts* the “compound” input type:

`\chcpd[⟨options⟩]{⟨compound⟩}`
 Typeset single compounds.

8 Special Input Types

There are some “special type” input groups.

8.1 Single Token Groups

The first kind are groups which consist of only one token, namely of the following ones:

`\ch{< + >}` +

Creates the plus sign between compounds with space around it:

`\ch{2 Na + Cl2}` 2 Na + Cl₂

`\ch{ v }↓`

Sign for precipitate: `\ch{BaSO4 v}` BaSO₄↓

`\ch{ ^ }↑`

Sign for escaping gas³: `\ch{H2 ^}` H₂↑

The space left and right of the plus sign can be set with this option:

`plus-space = {<skip>}`

Default: .3em plus .1em minus .1em

A rubber length.

`plus-penalty = {<num>}`

Default: 700

The penalty that is inserted after the plus sign for (dis-)allowing line breaks.


Introduced in
version 4.0a

| | |
|---|-------|
| 1 <code>\ch{A + B}\par</code> | A + B |
| 2 <code>\ch[plus-space=4pt]{A + B}</code> | A + B |

8.2 Option Input

Sometimes you might want to apply an option only to a part of a, say, reaction. Of course you have the possibility to use `\ch` several times.

| |
|--|
| 1 <code>\ch{H2O +}\textcolor{red}{\ch{H2SO4}}\ch{-> H3O+ + HSO4-} \par</code> |
| 2 <code>\ch{H2O +}\ch[subscript-vshift=2pt]{H2SO4}\ch{-> H3O+ + HSO4-}</code> |



This, however, interrupts the input in your source and *may* mess with the spacing. That's why there is an alternative:

`\ch @ {<options> }`



The options specified this way will be valid *only* until the next compound is set.

3. Is this the correct English term? Please correct me if it isn't.





```

1 \ch{H2O +}\textcolor{red}{\ch{H2SO4}}\ch{-> H3O+ + HSO4-} \par
2 \ch{H2O + @\format=\color{red}} H2SO4 -> H3O+ + HSO4-} \par
3 or of course:\par
4 \ch{H2O + \textcolor{red}{H2SO4} -> H3O+ + HSO4-}\par\bigskip
5 \ch{H2O +}\ch[subscript-vshift=2pt]{H2SO4}\ch{-> H3O+ + HSO4-} \par
6 \ch{H2O + @\subscript-vshift=2pt} H2SO4 -> H3O+ + HSO4-}

```



or of course:



9 Escaped Input

In some cases it may be desirable to prevent **CHEMFORMULA** from parsing the input. This can be done in two ways.

9.1 Text

If you put something between " " or ' ' then the input will be treated as normal text, except that spaces are not allowed and have to be input with ~.

`\ch{ "<escaped text>" }`

One of two possibilities to *escape* **CHEMFORMULA**'s parsing.


`\ch{ '<escaped text>' }`

The second of two possibilities to *escape* **CHEMFORMULA**'s parsing.

```

1 \ch{"\ox{2,Ca}" 0} \par
2 \ch{"\ldots\," Na + "\ldots\," Cl2 -> "\ldots\," NaCl} \par
3 \ch{'A~->~B'}

```



In many cases you won't need to escape the input. But when you get into trouble when using a command inside `\ch` try hiding it.

9.2 Math

If you especially want to input math you just enclose it with `$ $`. This output is different from the escaped text as it is followed by a space. The reasoning behind this is that I assume math will mostly be used to replace stoichiometric factors.

`\ch{ $\langle escaped math \rangle$ }`

One of two possibilities to *escape* **CHEMFORMULA**'s parsing into math mode.

`\ch{ \(\langle escaped math \rangle\) }`

The second of two possibilities to *escape* **CHEMFORMULA**'s parsing into math mode.

| | |
|---|--|
| 1 escaped text: <code>\ch{"\$x\$" H2O} \par</code> | escaped text: $x\text{H}_2\text{O}$ |
| 2 escaped math : <code>\ch{\$x\$ H2O} \par</code> | escaped math: $x\text{H}_2\text{O}$ |
| 3 also escaped math : <code>\ch{\(x\) H2O} \par</code> | also escaped math: $x\text{H}_2\text{O}$ |
| 4 <code>\ch{\$2n\$ Na + \$n\$ Cl2 -> \$2n\$ NaCl}</code> | $2n\text{Na} + n\text{Cl}_2 \longrightarrow 2n\text{NaCl}$ |

The space that is inserted after a math group can be edited:

`math-space = {\langle skip \rangle}`

Default: .1667em plus .0333em minus .0117em

A rubber length.

```
1 \ch{$2n$ Na + $n$ Cl2 -> $2n$ NaCl} \par
2 \chemsetup[chemformula]{math-space=.25em}
3 \ch{$2n$ Na + $n$ Cl2 -> $2n$ NaCl} \par
4 \ch{$A->B$}
```

$2n\text{Na} + n\text{Cl}_2 \longrightarrow 2n\text{NaCl}$
 $2n\text{Na} + n\text{Cl}_2 \longrightarrow 2n\text{NaCl}$
 $A- > B$

10 Arrows

10.1 Arrow types

Arrows are input in the same intuitive way they are with `mhchem`. There are various different types:

`\ch{ -> }` \longrightarrow

standard right arrow

`\ch{ <- }` \longleftarrow

standard left arrow

`\ch{ -/> }` \longrightarrow

does not react (right)

`\ch{ </- }` \longleftarrow

does not react (left)

`\ch{ <-> }` \longleftrightarrow

resonance arrow

`\ch{ <> }` \rightleftharpoons

reaction in both directions

`\ch{ == }` $=$

stoichiometric equation

`\ch{ <=> }` \rightleftharpoons

equilibrium arrow

`\ch{ <=>> }` \rightleftharpoons

unbalanced equilibrium arrow to the right

`\ch{ <<=> }` \rightleftharpoons

unbalanced equilibrium arrow to the left

`\ch{ <0> }` \longleftrightarrow








isolobal arrow

All these arrows are drawn with TikZ.

```

1 \ch{H2 + Cl2 -> 2 HCl} \par
2 \ch{H2O + CO3^2- <=> OH- + HCO3-} \par
3 \ch{A <- B} \par
4 \ch{\{[CH2=CH-CH2]- <-> \}[CH2-CH=CH2]- \}} \par
5 \ch{A <> B} \par
6 \ch{H+ + OH- <=>> H2O} \par
7 \ch{2 NO2 <<=> N2O4}

```




10.2 Labels

The arrows take two optional arguments to label them.

`\ch{ ->[<above>][<below>] }`


Add text above or under an arrow.

```
1 \ch{A ->[a] B} \par
2 \ch{A ->[a][b] B} \par
3 \ch{A ->[\SI{100}{\celsius}] B}
```




The label text can be parsed separately from the arrow. The recipe is easy: leave blanks.

```
1 \ch{A ->[H2O] B} \par
2 \ch{A ->[ H2O ] B} \par
3 \ch{A ->[ "\ox{2,Ca}" F2 ] B} \par
4 \ch{A ->[$\Delta$,~ \[H+ \]] B}
```



If you leave the blanks **CHEMFORMULA** treats the groups inside the square brackets as separated input types. The arrow reads its arguments *afterwards*. As you can see the arrows “grow” with the length of the labels. What stays constant is the part that protrudes the labels. As you also can see in the last example square brackets inside the arrow arguments should be produced using `\[` and `\]`. They keep their usual meaning outside `\ch`. These commands are necessary since the usual grouping (*i. e.*, hiding the brackets inside curly brackets) doesn’t work due to the way `\ch` reads its argument.

```
1 \ch{A ->[a] B} \par
2 \ch{A ->[ab] B} \par
3 \ch{A ->[abc] B} \par
4 \ch{A ->[abc~abc] B} \par
5 % needs the 'chemfig' package:
6 \setatomsep{15pt}
7 \ch{A ->[ "\chemfig{-[:30]-[:-30]OH}" ] B} \par
8 \ch{A ->[[ ]] B} vs. \ch{A ->[\[ \]] B}
```



10.3 Customization

These are the options which enable you to customize the arrows:

`arrow-offset = {⟨dim⟩}` Default: .75em

This is the length that an arrow protrudes a label on both sides. This means an empty arrow's length is two times `arrow-offset`.

`arrow-min-length = {⟨dim⟩}` Default: 0pt

Introduced in
version 3.6b

The minimal length an error must have unless two times `arrow-offset = {⟨p⟩}` plus the width of the label is larger.

`arrow-yshift = {⟨dim⟩}` Default: 0pt

Shifts an arrow up (positive value) or down (negative value).

`arrow-ratio = {⟨factor⟩}` Default: .6

The ratio of the arrow lengths of the unbalanced equilibrium. .4 would mean that the length of the shorter arrow is 0.4× the length of the longer arrow.

`compound-sep = {⟨dim⟩}` Default: .5em

The space between compounds and the arrows.

`label-offset = {⟨dim⟩}` Default: 2pt

The space between the labels and the arrows.

`label-style = {⟨font command⟩}` Default: `\footnotesize`

The relative font size of the labels.

`arrow-penalty = {⟨num⟩}` Default: 0

Introduced in
version 4.0a

The penalty that is inserted after an arrow for (dis-)allowing line breaks.

`arrow-style = {⟨TikZ⟩}` (initially empty)

Introduced in
version 4.1a

Additional TikZ keys for formatting the arrows.

The following code shows the effect of the different options on the `<=>` arrow:

```

1 standard: \ch{A <=>[x][y] B} \par
2 longer: \ch[arrow-offset=12pt]{A <=>[x][y] B} \par
3 higher: \ch[arrow-yshift=2pt]{A <=>[x][y] B} \par
4 more balanced: \ch[arrow-ratio=.8]{A <=>[x][y] B} \par
5 labels further away: \ch[label-offset=4pt]{A <=>[x][y] B} \par
6 larger distance to compounds: \ch[compound-sep=2ex]{A <=>[x][y] B} \par
7 smaller labels: \ch[label-style=\tiny]{A <=>[x][y] B}

```

standard: A $\xrightarrow{\frac{x}{y}}$ B
 longer: A $\xrightarrow{\frac{x}{y}}$ B
 higher: A $\xrightarrow{\frac{x}{y}}$ B

more balanced: $A \xrightleftharpoons[x]{x} B$
 labels further away: $A \xrightleftharpoons[y]{x} B$
 larger distance to compounds: $A \xrightleftharpoons[y]{x} B$
 smaller labels: $A \xrightleftharpoons[y]{x} B$

10.4 Modify Arrow Types

The arrows are defined with the command

`\DeclareChemArrow{<type>}{<TikZ>}`

Define a new arrow type *<type>*.

<type> is the sequence of tokens that is replaced with the actual arrow code. For example the basic arrow is defined via

```
1 \DeclareChemArrow{->}{
2   \draw[chemarrow,-cf] (cf_arrow_start) -- (cf_arrow_end) ;
3 }
```

In order to define arrows yourself you need to know the basics of TikZ.⁴ The predefined arrows use the arrow tips `cf`, `left cf` and `right cf`. They also all except the net reaction arrow `==` use the TikZ-style `chemarrow` that you should use, too, if you want the option `arrow-style` to have an effect.

There are some predefined coordinates you can and should use. For completeness' sake the arrow tips and the TikZ-style are also listed:

`(cf_arrow_start)`

The beginning of the arrow.

`(cf_arrow_end)`

The end of the arrow.

`(cf_arrow_mid)`

The mid of the arrow.

`(cf_arrow_mid_start)`

The beginning of the shorter arrow in types like `<=>`.

`(cf_arrow_mid_end)`

The end of the shorter arrow in types like `<=>`.

`cf`

A double-sided arrow tip.

⁴. Please see the pgfmanual for details.

`left cf`

A left-sided arrow tip.

`right cf`

A right-sided arrow tip.


`chemarrow`

CHEMFORMULA's TikZ-style that is applied to the arrows and set with `arrow-style`

```

1 \DeclareChemArrow{.>}{
2   \draw[chemarrow,-cf,dotted,red] (cf_arrow_start) -- (cf_arrow_end);
3 }
4 \DeclareChemArrow{n>}{
5   \draw[chemarrow,-cf]
6     (cf_arrow_start)
7     .. controls ([yshift=3ex]cf_arrow_mid) ..
8     (cf_arrow_end);
9 }
10 \ch{A .> B} \ch{A .>[a][b] B} \ch{A n> B}

```



If you want to redefine an existing arrow there are two commands you can use:

`\RenewChemArrow{<type>}{<TikZ>}`

Redefine the arrow type `<type>`.

`\ShowChemArrow{<type>}`


Print out the current definition of the arrow type `<type>`.

The second one gives the current definition, the first one redefines an existing arrow.

```

1 \texttt{\ShowChemArrow{->}} \par
2 \RenewChemArrow{->}{\draw[chemarrow,->,red] (cf_arrow_start) -- (cf_arrow_end)
3   ;}
4 \texttt{\ShowChemArrow{->}} \par
5 \ch{A -> B}

```



10.5 Standalone Arrows

Introduced in
version 4.0

CHEMFORMULA offers a command that *only accepts* the “arrow” input type:

`\charrow{<type>}[<above>][<below>]`

Print the arrow type <type>.

This command is internally used for the arrows, too, when `\ch` is parsed.

11 Names

11.1 Syntax

CHEMFORMULA has a built-in syntax to write text under a compound. In a way it works very similar to the arrows.

`\ch{ !(<text>)(<formula>) }`

Writes <text> below <formula>.

If an exclamation mark is followed by a pair of parentheses **CHEMFORMULA** will parse it this way:

```
1 \ch{!(ethanol)( CH2CH2OH )}
```

CH₂CH₂OH
ethanol

The same what's true for the arrows arguments holds for these arguments: if you leave blanks the different parts will be treated according to their input type before the text is set below the formula.

```
1 \ch{!(water)(H2O)} \quad
2 \ch{!( "\textcolor{blue}{water}" )( H2O )} \quad
3 \ch{!( $2n-1$ )( H2O )} \quad
4 \ch{!( H2O )( H2O )} \quad
5 \ch{!(oxonium)( H3O+ )}
```

| | | | | |
|------------------|------------------|------------------|------------------|-------------------------------|
| H ₂ O | H ₂ O | H ₂ O | H ₂ O | H ₃ O ⁺ |
| water | water | 2n - 1 | H ₂ O | oxonium |

If for some reason you want to insert an exclamation mark *without* it creating a name you only have to make sure it isn't followed by parentheses.

```
1 \ch{H2O~(!)} \quad
2 \ch{A!{ }{ }{ }}
```

H₂O (!)
A!()

11.2 Customization

CHEMFORMULA provides two options to customize the output of the names:

name-format = {<commands>} Default: `\scriptsize\centering`

The format of the name. This can be arbitrary input.

name-width = <dim>|auto Default: auto

The width of the box where the label is put into. auto will detect the width of the name and set the box to this width.

```

1 \ch{!(acid)( H2SO4 ) -> B} \par
2 \ch[name-format=\sffamily\small]{!(acid)( H2SO4 ) -> B} \par
3 \ch[name-format=\scriptsize N:~]{!(acid)( H2SO4 ) -> B} \par
4 \ch[name-width=3em,name-format=\scriptsize\raggedright]{!(acid)( H2SO4 ) -> B}

```

$\text{H}_2\text{SO}_4 \longrightarrow \text{B}$
 acid
 $\text{H}_2\text{SO}_4 \longrightarrow \text{B}$
 acid
 $\text{H}_2\text{SO}_4 \longrightarrow \text{B}$
 N: acid
 $\text{H}_2\text{SO}_4 \longrightarrow \text{B}$
 acid

11.3 Standalone Names

Introduced in
version 4.0

CHEMFORMULA offers a command that allows the usage of the “name” syntax in normal text. This is the command that a bang is replaced with in **CHEMFORMULA**’s formulas, actually. Both arguments are mandatory.

`\chname(<text 1>)(<text 2>)`

The command that is used internally for placing <text 1> below of <text 2>.


12 Format and Font

In the standard setting **CHEMFORMULA** doesn’t make any default changes to the font of the formula output. Let’s take a look at a nonsense input which shows all features:

```

1 \newcommand*\sample{%
2   \ch{H2C-C+C-CH=CH+ + CrO4^2-
3     <=>[x][y]
4     2.5 Cl^{-.} + 3_1/2 Na*OH_{(aq)} + !(name)( A^n ) "\LaTeXe"}
5 }
6 \sample

```




Now we're going to change different aspects of the font a look what happens:

```

1 \sffamily Hallo \sample \
2 \ttfamily Hallo \sample \normalfont \
3 \bfseries Hallo \sample \normalfont \
4 \itshape Hallo \sample

```



As you can see most features adapt to the surrounding font.

If you want to change the default format you need to use this option:


`format = {\langle anything \rangle}` (initially empty)

Adds arbitrary code before the output of `\ch`.

```

1 \definecolor{newblue}{rgb}{.1,.1,.5}
2 \chemsetup[chemformula]{format=\color{newblue}\sffamily}
3 \sffamily Hallo \sample \
4 \ttfamily Hallo \sample \normalfont \
5 \bfseries Hallo \sample \normalfont \
6 \itshape Hallo \sample

```



You can also specifically change the fontfamily, fontseries and fontshape of the output.

font-family = {<family>} (initially empty)

Changes the fontfamily of the output with `\fontfamily{<family>}`.

font-series = {<series>} (initially empty)

Changes the fontseries of the output with `\fontseries{<series>}`.

font-shape = {<shape>} (initially empty)

Changes the fontshape of the output with `\fontshape{<shape>}`.

```

1 \chemsetup[chemformula]{font-series=bx}
2 Hallo \sample \par
3 \sffamily Hallo \sample \normalfont \par
4 \chemsetup[chemformula]{font-family=lmss,font-series=m} Hallo \sample
5 \normalfont \par
6 \itshape Hallo \sample

```

Hallo $\text{H}_2\text{C}-\text{C}\equiv\text{C}-\text{CH}=\text{CH}^+ + \text{CrO}_4^{2-} \xrightleftharpoons[\text{y}]{\text{x}} 2.5 \text{Cl}^- + 3\frac{1}{2} \text{Na} \cdot \text{OH}_{(\text{aq})} + \overset{\text{name}}{\text{A}^n \text{L}^{\text{A}}\text{T}_{\text{E}}\text{X} 2_{\text{E}}}$

Hallo $\text{H}_2\text{C}-\text{C}\equiv\text{C}-\text{CH}=\text{CH}^+ + \text{CrO}_4^{2-} \xrightleftharpoons[\text{y}]{\text{x}} 2.5 \text{Cl}^- + 3\frac{1}{2} \text{Na} \cdot \text{OH}_{(\text{aq})} + \overset{\text{name}}{\text{A}^n \text{L}^{\text{A}}\text{T}_{\text{E}}\text{X} 2_{\text{E}}}$

Hallo $\text{H}_2\text{C}-\text{C}\equiv\text{C}-\text{CH}=\text{CH}^+ + \text{CrO}_4^{2-} \xrightleftharpoons[\text{y}]{\text{x}} 2.5 \text{Cl}^- + 3\frac{1}{2} \text{Na} \cdot \text{OH}_{(\text{aq})} + \overset{\text{name}}{\text{A}^n \text{L}^{\text{A}}\text{T}_{\text{E}}\text{X} 2_{\text{E}}}$

Hallo $\text{H}_2\text{C}-\text{C}\equiv\text{C}-\text{CH}=\text{CH}^+ + \text{CrO}_4^{2-} \xrightleftharpoons[\text{y}]{\text{x}} 2.5 \text{Cl}^- + 3\frac{1}{2} \text{Na} \cdot \text{OH}_{(\text{aq})} + \overset{\text{name}}{\text{A}^n \text{L}^{\text{A}}\text{T}_{\text{E}}\text{X} 2_{\text{E}}}$

If you're using $\text{X}_{\text{Y}}\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}$ or $\text{LuaL}^{\text{A}}\text{T}_{\text{E}}\text{X}$ and have loaded `fontspec` you have the possibility to set the font with it:

font-spec = {} (initially empty)

Use font for **CHEMFORMULA**'s formulas.

or with options

font-spec = {[<options>]}

Use font with options <options> for **CHEMFORMULA**'s formulas.

Since this document is typeset with $\text{pdfL}^{\text{A}}\text{T}_{\text{E}}\text{X}$ the option cannot be demonstrated here.


13 Usage In Math Equations

The `\ch` command can be used inside math equations. It recognizes `\\` and `&` and passes them on. However, you can't use the optional arguments of `\\` inside `\ch`.

```

1 \begin{align}
2 \ch{
3   H2O & \rightarrow[a] H2SO4 \\
4   Cl2 & \rightarrow[x][y] CH4
5 }
6 \end{align}
7 \begin{align*}
8 \ch{
9   RNO2 & \rightleftharpoons[+ e^-]{ } RNO2^{\cdot-} \\
10  RNO2^{\cdot-} & \rightleftharpoons[+ e^-]{ } RNO2^{2-}
11 }
12 \end{align*}

```



14 Usage with TikZ or pgfplots and externalization

Introduced in
version 4.1

Since **CHEMFORMULA** uses TikZ to draw reaction arrows and bonds they would be externalized, too, if you use that facility with TikZ or pgfplots⁵ [Feu13]. This may not be desirable since they are very small pictures maybe containing of a single line. This is why **CHEMFORMULA**'s default behaviour is to disable externalization for it's bonds and arrows. This can be turned on and off through the following option:

tikz-external-disable = true|false Default: true
dis- or enable TikZ' externalization mechanism for **CHEMFORMULA**'s arrows and bonds.

If you should be using a formula that contains bonds or arrows inside of a tikzpicture that is externalized you should locally enable it for **CHEMFORMULA**, too:

```

1 \begin{tikzpicture}
2   \setchemformula{tikz-external-disable=false}
3   \begin{axis}[xlabel={\ch{2 H+ + 2 e- -> H2}}]
4     \addplot ... ;
5   \end{axis}

```

5. on CTAN as pgfplots: <http://mirrors.ctan.org/macros/latex/contrib/pgfplots/>

```
6 \end{tikzpicture}
```

15 Lewis Formulae

Introduced in
version 4.2

CHEMFORMULA offers a command to typeset Lewis formulae. This does not mean Lewis structures! Those can be achieved using the chemfig package [Tel12]. **CHEMFORMULA** provides the possibility to draw electrons as dots and pairs of dots or a line around an atom.

`\chlewis[⟨options⟩]{⟨electron spec⟩}{⟨atom⟩}`

Draws electrons around the `⟨atom⟩` according to `⟨electron spec⟩`.

Electrons are specified by the angle to the horizontal in the counter-clockwise direction. The default appearance is a pair of electrons drawn as a pair of dots. Other specifications can be chosen. The specification follows the pattern `⟨angle⟩⟨separator⟩`. `⟨angle⟩` is a positive or negative integer denoting the angle counter clockwise to the horizontal where the electrons should be drawn. `⟨separator⟩` is either a dot (., single electron), a colon (:, electron pair), a vertical line (|, electron pair), an o (o, empty pair), or a comma (, default spec).

`\chlewis{⟨angle1⟩⟨type1⟩⟨angle2⟩⟨type2⟩}{⟨atom⟩}`

For example: `\chlewis{0,180}{O}` gives $\text{:}\ddot{\text{O}}\text{:}$ and `\chlewis{0.90.180.270.}{C}` gives $\cdot\dot{\text{C}}\cdot$.

The appearance can be influenced by a number of options:

`lewis-default = .|:|o|single|pair|pair (dotted)|pair (line)|empty` Default: pair
Sets the default type that is used when no type is given in `⟨electron spec⟩`.

`lewis-distance = {⟨dim⟩}` Default: 1ex
The distance of two electrons in a pair.

`lewis-line-length = {⟨dim⟩}` Default: 1.5ex
The length of the line representing an electron pair.

`lewis-line-width = {⟨dim⟩}` Default: 1pt
The thickness of a line representing an electron pair.

`lewis-offset = {⟨dim⟩}` Default: .5ex
The distance of the symbols from the atom.

The dots are drawn according to the `radical-radius` option mentioned in section 7.6.

The basic usage should be more or less self-explaining:

```
1 \chlewis{0:90|180.270}{O}
2 \quad
3 \chlewis{45,135}{O}
4 \quad
5 \chlewis{0o}{Na}
```



The next example shows the effect of some of the options:

16 Further Examples

```

1 \chlewis[lewis-default=.]{23,68,113,158,203,248,293,338}{X}
2 \quad
3 \chlewis{0,90,180,270}{X}
4 \quad
5 \chlewis[lewis-distance=1.25ex]{0,90,180,270}{X}
6 \quad
7 \chlewis[lewis-distance=.75ex,radical-radius=.5pt]{0,90,180,270}{X}
8 \quad
9 \chlewis[
10   radical-radius=.5pt,
11   lewis-default=.
12 ]{23,68,113,158,203,248,293,338}{X}


```



```

1 \ch{
2   !($1s^22s^1$)( "\chlewis{180.}{Li}" ) +
3   !($1s^22s^22p^5$)( "\chlewis{0.90,180,270}{F}" )
4   ->
5   !($1s^2$)( Li+ ) + !($1s^22s^22p^6$)( "\chlewis{0,90,180,270}{F}" {}- )
6 }

```




16 Further Examples

This section presents some examples of a possible usage.

```

1 \begin{reaction}[Synthese von Alkanen]
2   !(Synthesegas)( $n$ CO + $(2n+1)$ H2 )
3   ->[\SI{200}{\celsius}][\CoNi]
4   C_{n$}H_{2n+2$} + $n$ H2O
5 \end{reaction}

```




16 Further Examples

```

1 \begin{reactions*}
2   "a)" && CH4 + Cl2 &-> CH3Cl + HCl && "{\small Chlormethan/Methylchlorid}"
3   "b)" && CH3Cl + Cl2 &-> CH2Cl2 + HCl && "{\small Dichlormethan/
4   Methylenchlorid}" \\
5   "c)" && CH2Cl2 + Cl2 &-> CHCl3 + HCl && "{\small Trichlormethan/Chloroform}"
6   "d)" && CHCl3 + Cl2 &-> CCl4 + HCl && "{\small Tetrachlormethan/
7   Tetrachlorkohlenstoff}"
8 \end{reactions*}


```



```

1 \chemsetup[ox]{parse=false}
2 \ch{
3   "\ox{\delm,C}" -{ } "\ox{\delp,M}" \quad
4   ( <-> "\ox{\delp,C}" -{ } "\ox{\delm,Br}" )
5 } \\
6 \ch[adduct-space=0pt]{X. + .Y <=> X-Y + Bindungsenergie} \\
7 \ch[name-format=\normalsize]{
8   !(\State{H}{f}\quad) !(\textcolor{red}{??})( CH4\gas{ } ) +
9   !(\num{0})( 2 O2\gas{ } )
10  ->
11   !(\num{-94.3})( CO2\gas{ } ) +
12   !(\num{-57.9})( H2O\lqd{ } ) +
13   !(\num{-192.1})( "\State{H}" )
14 }

```



```

1 \newcommand\cstack[1]{%
2   \clap{%


```

16 Further Examples

```

3   \begin{tabular}{c}
4     #1
5   \end{tabular}
6   }%
7 }
8 \begin{reactions*}
9   CH3MgBr + "\ox*{1,Cu}" X &
10  -> "\glqq" CH3 "\ox*{1,Cu}\grqq" + MgBrX "\quad X~$=$~Br,I,CN" \\\
11  2 MeLi + CuI &
12  -> !(\cstack{Dimethylcuprat\\(Gilman-Cuprat)})( Me2CuLi ) + Li
13 \end{reactions*}


```



```

1 % needs 'chemfig'
2 \begin{reactions*}
3   H3C-CH3 + Cl2 & &
4   ->[\Delta][h\nu] H3CCH2Cl + HCl & &
5   "\Enthalpy{-27.1}" \\\
6   H3C-CH3 + "\Lewis{0.,Cl}" & &
7   -> H3CCH2 "\Lewis{0.,\vphantom{H}}" + HCl & &
8   "\Enthalpy{-5.0}" \\\
9   H3C-CH2 "\Lewis{0.,\vphantom{H}}" + Cl2 & &
10  -> H3CCH2Cl + "\Lewis{0.,Cl}" & &
11  "\Enthalpy{-23.0}"
12 \end{reactions*}

```



The following example shows how the cancelling of compounds could be done.⁶


6. Inspired by a question on TeX.SE: <http://tex.stackexchange.com/q/30118/5049>

References

```

1 % needs 'cancel'
2 \begin{align*}
3   \ch{\cancel{HCOOH}\aq} + H2O\lqd{} &\rightleftharpoons H3O^+\aq{} + \cancel{HCOO^-\aq{}} \\
4   \ch{\cancel{HCOO^-\aq{}} + H2O\lqd{}} &\rightleftharpoons \cancel{HCOOH\aq} + OH^-\aq{} \\
5   \cline{1-2}
6   \ch{H2O\lqd{}} + H2O\lqd{} &\rightleftharpoons H3O^+\aq{} + OH^-\aq{}
7 \end{align*}

```



References

- [Ame02] American Mathematical Society. amsmath. version 2.0, Feb. 25, 2002.
URL: <http://mirror.ctan.org/macros/latex/required/amsmath/>.
- [Coh+08] E. Richard Cohan et al.
“Quantities, Symbols and Units in Physical Chemistry”, *IUPAC Green Book*.
3rd Edition. 2nd Printing. IUPAC & RSC Publishing, Cambridge, 2008.
- [Feu13] Christian Feuersänger. pgfplots. version 1.9, Oct. 4, 2013.
URL: <http://mirror.ctan.org/graphics/pgf/contrib/pgfplots/>.
- [KN12] Markus Kohm and Frank Neukahm. KOMA-Script. version 3.11b, July 29, 2012.
URL: <http://mirror.ctan.org/macros/latex/contrib/koma-script/>.
- [Rei98] Axel Reichert. nicefrac. version 0.9b, Aug. 4, 1998.
URL: <http://mirror.ctan.org/macros/latex/contrib/units/>.
- [Tan10] Till Tantau. TikZ/pgf. version 2.10, Oct. 25, 2010.
URL: <http://mirror.ctan.org/graphics/pgf/>.
- [Tel12] Christian Tellechea. chemfig. version 1.0g, Nov. 16, 2012.
URL: <http://mirror.ctan.org/macros/generic/chemfig/>.
- [The13a] The L^AT_EX3 Project Team. l3kernel. version SVN 4582, July 28, 2013.
URL: <http://mirror.ctan.org/macros/latex/contrib/l3kernel/>.
- [The13b] The L^AT_EX3 Project Team. l3packages. version SVN 4582, July 28, 2013.
URL: <http://mirror.ctan.org/macros/latex/contrib/l3packages/>.

Index

A

`adduct-space` 12, 15
 American Mathematical Society 3
`amsmath` (package) 3
`\aq` 4, 8, 25, 33
`arrow-min-length` 21
`arrow-offset` 21
`arrow-penalty` 21
`arrow-ratio` 21
`arrow-style` 2, 21 ff.
`arrow-yshift` 21

B

`\bond` 10 ff., 15
`bond-length` 12
`bond-offset` 13
`bond-penalty` 13
`bond-style` 13

C

`\ch` 3–10, 12–21, 23–28, 30 ff., 33
`charge-hshift` 10, 12 f.
`charge-style` 12
`charge-vshift` 12
`\charrow` 24
`\chcpd` 15
`chemfig` (package) 29
`chemformula` 3 f.
`\chemformula@bondlength` 12
`\chemsetup` 3, 5, 9, 12–15, 18, 26 f., 31
`\chlewis` 2, 29 f.
`\chname` 25
 Cohan, E. Richard 5, 10, 13
`compound-sep` 21
 CTAN 3, 28

D

`decimal-marker` 6
`\DeclareChemArrow` 22 f.
`\DeclareChemBond` 11 f.
`\DeclareChemBondAlias` 11
`\delm` 31
`\delp` 31
`\Delta` 20, 32

E

`\Enthalpy` 32

F

Feuersänger, Christian 28
`font-family` 27

`font-series` 27
`font-shape` 27
`font-spec` 27
`fontspec` (package) 27
`format` 26
`frac-math-cmd` 6
`frac-style` 6 f.

G

`\gas` 31

H

Hensel, Martin 2

K

Kohm, Markus 3
 KOMA-Script (bundle) 3

L

`l3kernel` (bundle) 3
`l3keys2e` (package) 3
`l3packages` (bundle) 3
`label-offset` 21
`label-style` 21
`lewis-default` 29
`lewis-distance` 29
`lewis-line-length` 29
`lewis-line-width` 29
`lewis-offset` 29
 LPPL 3
`\lqd` 31, 33

M

`math-space` 18
`\mch` 9
`mhchem` (package) 2 f., 18

N

`name-format` 25
`name-width` 25
 Neukahm, Frank 3
`nicefrac` (package) 3, 7
 Niederberger, Clemens 1

O

`\ox` 8, 17, 20, 31 f.

P

`pgfmanual` (manual) 22
`pgfplots` (package) 28
`plus-penalty` 16
`plus-space` 16

INDEX

Q

“Quantities, Symbols and Units in Physical Chemistry”,
IUPAC Green Book 5, 10, 13

R

`radical-hshift` 13
`radical-radius` 13, 29
`radical-space` 13
`radical-style` 13
`radical-vshift` 13
Reichert, Axel 3
`\RenewChemArrow` 23
`\RenewChemBond` 11

S

`scrfile` (package) 3
`\setchemformula` 3, 5, 28
`\ShowChemArrow` 23

`\ShowChemBond` 11
`\State` 31
`stoich-paren-parse` 6
`stoich-space` 6 f.
`subscript-style` 12
`subscript-vshift` 12, 14

T

Tantau, Till 3
Tellechea, Christian 29
The L^AT_EX3 Project Team 3
`tikz` (package) 3
`tikz-external-disable` 28
TikZ/pgf (package) 3

X

`xfrac` (package) 3, 6
`xparse` (package) 3